

this week

(i)

Environmental Policies >

> Land Management





- Marine conservation
- Conservation status

RELATED TERMS

- > Penguin
- > Fishery
- > Marine biology
- > Antarctic krill
- > Zoo

current and proposed MPAs. The study, which tracked eight penguins with satellite tags over a year, also found that they commonly traveled over 1,200 kilometers (745 miles) beyond the species range defined by the International Union for Conservation of Nature (IUCN), which is based on studies of adult emperor penguins from a few other colonies. Considered immature until about 4 years of age, juvenile emperor penguins are more vulnerable than adults because they have not fully developed foraging and predator avoidance skills. As climate change reduces sea-ice habitat and opens up new areas of the Southern Ocean to commercial fishing, the researchers conclude that greatly expanded MPAs are crucial to protect this iconic, yetthreatened, penguin species at every life stage.

"While everyone is looking at the adult population, the juvenile population -- which leaves the relative safety of its parents at about five months -- is neither monitored nor protected," said Dan Zitterbart, a WHOI associate scientist. "The current and proposed MPAs in the Southern Ocean only include the range of adult emperor penguins, which do not travel as far as juveniles. From a conservation perspective, it's important to know where these juveniles go. It's one more piece of the puzzle to protect their marine habitat."

"Emperor penguins have such low fecundity, if you do not protect juveniles, they may not ever become breeding adults," he continued.

Zitterbart and colleagues at the Centre Scientifique de Monaco (CSM), Centre National de la Recherche Scientifique (CNRS) and Université de Strasbourg in France, and the Alfred-Wegener Institute (AWI) in Germany are conducting a long-term monitoring study of the Atka Bay emperor penguin colony near Neumayer Station III, on the Weddell Sea. The Weddell Sea area is home to one-third of established emperor penguin colonies, and research shows that colonies in the region, including the Ross Sea, are less vulnerable to climate-induced melting than other areas of the Antarctic.

"Some of the Weddell Sea colonies are expected to still be present 50 to 100 years from now," said Aymeric Houstin, a WHOI post-doctoral investigator and the lead author of the study. "It's important to preserve colonies that will be able to endure climate change, as they could become a refuge for the entire population of emperor penguins."

According to studies, 12 percent of the area under CCAMLR jurisdiction is currently protected as an MPA, and less than 5 percent is considered a "no-take" area. For several years, the 26 members of CCAMLR have been considering three new MPAs in the region, including the Weddell Sea MPA, first developed by Germany, and submitted by the European Union in 2013. While this MPA would cover an area of 2.2 million square kilometers (0.85 million square miles), preserving one of the most pristine ecosystems in the world and a critical zone for global ocean circulation, the authors say that the boundaries are inadequate to protect juvenile emperor penguins.

PLANTS & ANIMALS

New Study Links Ultra-Processed Foods and Colorectal Cancer in Men

Cravings for Fatty Foods Traced to Gut-Brain Connection

Dolphins Form Largest Alliance Network Outside Humans, Study Finds

EARTH & CLIMATE

Faster in the Past: New Seafloor Images of West Antarctic Ice Sheet Upend Understanding of Thwaites Glacier Retreat

People Generate Their Own Oxidation Field and Change the Indoor Air Chemistry Around Them

Risk of Multiple Climate Tipping Points Escalates Above 1.5°C Global Warming

FOSSILS & RUINS

Modern Humans Generate More Brain Neurons **Than Neanderthals**

Stone Age Surgery: Earliest Evidence of **Amputation Found**

DNA in Viking Feces Sheds New Light on 55,000-Year-Old Relationship Between Gut Companions

ADVERTISEMENT

Advertisements



> Fish farming

"The Weddell Sea MPA design, as the other MPAs around Antarctica, should include the distribution at sea of all age-classes of the emperor penguin population -- not only the adults from a few study colonies," said Céline Le Bohec, of CNRS/Université de Strasbourg France and the Centre Scientifique de Monaco. "Juveniles are currently clearly lacking protection and their presence in the Northern waters needs to be considered in the future, especially regarding the development of fisheries in those regions."

Over the next decades, the researchers plan to continue tagging both adult and juvenile penguins from the Atka Bay colony to track their movements and behavior as the environment changes. With more long-term data, Houstin suggests that a "dynamic MPA" could be developed with shifting boundaries, based on predictions of penguins' movements throughout the year.

"This notion of a dynamic network of MPAs is really essential," Le Bohec said. "It's certainly the way to continue the dialogue with the fishing industry to ensure the resource is used in a sustainable manner, to ultimately preserve the unique biodiversity of these sensitive polar regions."

Story Source:

Materials provided by Woods Hole Oceanographic Institution. Note: Content may be edited for style and length.

Journal Reference:

1. Aymeric Houstin, Daniel P. Zitterbart, Karine Heerah, Olaf Eisen, Víctor Planas-Bielsa, Ben Fabry, Céline Le Bohec. Juvenile emperor penguin range calls for extended conservation measures in the Southern Ocean. Royal Society Open Science, 2022; 9 (8) DOI: 10.1098/rsos.211708

Cite This Page:

APA Chicago

Woods Hole Oceanographic Institution. "Marine Protected Areas in Antarctica should include young emperor penguins, scientists say." ScienceDaily. ScienceDaily, 31 August 2022. <www.sciencedaily.com/releases/2022/08/220831111120.htm>.

MLA

Strange & Offbeat

PLANTS & ANIMALS

This Fungus Shrinks in Size to Better Infect the Brain

The Gene to Which We Owe Our Big Brain

What Killed Dinosaurs and Other Life on Earth?

EARTH & CLIMATE

Mysterious Diamonds Came from Outer Space

Changes in the Tree Canopy Facilitated the **Evolution of the First-Ever Gliding Reptile**

Study Unearths Ancient Reef Structure High and Dry on the Nullarbor Plain

FOSSILS & RUINS

Stone Age Surgery: Earliest Evidence of Amputation Found

Modern Humans Generate More Brain Neurons Than Neanderthals

These Female Hummingbirds Evolved to Look Like Males -- Apparently to Evade Aggression

Explore More

RELATED STORIES

from ScienceDaily

ADVERTISEMENT

Study Shows How Network of Marine Protected Areas Could Help **Safeguard Antarctic Penguins**

Jan. 20, 2021 — New research highlights how a proposed network of marine protected areas could help safeguard some of the most important areas at sea for breeding Antarctic ...

Marine Ecosystems: No Refuge from the Heat

Dec. 7, 2020 — Over the past several decades, marine protected areas (MPAs) have emerged as a favored conservation tool. By protecting marine species and safeguarding habitat, these reserves help buffer ecosystems ...

Fish Species Benefit from Marine Protection to Varying Extents

Jan. 8, 2020 — Marine protected areas reduce fish mortality by limiting harvesting and reducing habitat destruction. They are often designed and implemented to promote biodiversity conservation and sustainable ...

Climate Change Threatens Marine Protected Areas

May 8, 2018 — New research found that most marine life in Marine Protected Areas will not be able to tolerate warming ocean temperatures caused by greenhouse gas emissions. The study found that with continued ...



Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

Email Newsletters

RSS Feeds

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

f Facebook

Twitter

in LinkedIn

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

- Leave Feedback
- Contact Us

About This Site | Staff | Reviews | Contribute | Advertise | Privacy Policy | Editorial Policy | Terms of Use

Copyright 1995-2022 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners. Financial support for ScienceDaily comes from advertisements and referral programs, where indicated. CCPA: Do Not Sell My Information — GDPR: Privacy Settings —